

**PROTOCOLS** 

**MEDICAL** 

**EVIDENCE** 

**COVID RESOURCES** 

**EDUCATION & EVENTS** 

NEWS & **WEBINARS** 

**ABOUT FLCCC** 

DONATE

#### MORE RECOVERY PROTOCOLS

I-RECOVER: Long COVID Treatment

#### **TAKE ME TO:**

- Prevention
- Treatment

Recovery

Back to All Protocols

#### **SUPPORTING DOCUMENTS**

- Efficacy of Ivermectin
- Nutritional Therapeutics and
- Vitamins and Nutraceuticals **During Pregnancy**
- Nutrient Guide
- Traducciones de Nuestros Protocolos en Español
- Frequently Asked Questions

CC BY-NC-ND 4.0 (cc) (†) (\$) (=)

This work is licensed under

# I-RECOVER

### POST-VACCINE TREATMENT PROTOCOL

Download I-RECOVER Post-Vaccine Summary

Download I-RECOVER Post-Vaccine Protocol

Post-vaccine syndrome is a complex disease. Treatment must be individualized according to each patient's presenting symptoms and disease syndromes. Not all patients respond equally to the same intervention. Early treatment is essential; the response to treatment will likely be weaker when treatment is delayed.

This document is designed for healthcare providers caring for patients with symptoms following a COVID injection. While a handful of the therapies can be self-administered, we strongly recommend that patients consult with a healthcare provider before beginning any new treatment. (To find a provider, consult FLCCC's provider directory.)

There are also some important cautions and contraindications that should be carefully reviewed within the more comprehensive and detailed document called "An Approach to Managing Post-Vaccine Syndrome" and which should be discussed with a qualified provider as well.

This information is not intended to serve as a substitute for diagnosis, treatment, or advice from a qualified, licensed medical professional. The facts presented are offered as information — not medical advice. Any treatment protocol should be discussed with a trusted, licensed medical professional. Never stop or change medications without consulting your healthcare provider.

# **Treatment approach**

It is important to emphasize that, since there are no published reports detailing how to treat vaccine-injured patients, our treatment approach is based on the postulated pathogenetic mechanisms, principles of pharmacology, clinical observations, and feedback from vaccineinjured patients themselves. We are constantly updating the approach as new data emerges and based on consultation with trusted healthcare providers.

Patients with post-vaccine syndrome must not receive further COVID-19 vaccines of any type. Likewise, patients with long COVID should avoid all COVID vaccinations.

Patients with post-vaccine syndrome should do whatever they can to prevent themselves from getting COVID-19. This may include a preventative protocol (see <u>I-PREVENT</u>) or early treatment in the event you do contract the virus or suspect infection (see <u>I-CARE</u>). COVID-19 will likely exacerbate the symptoms of vaccine injury.

Once a patient has shown improvement, the various interventions should be reduced or stopped one at a time. A less intensive maintenance approach is then suggested.

The core problem in post-vaccine syndrome is long-lasting "immune dysregulation." The most important treatment goal is to help the body restore a healthy immune system — in other words, to let the body heal itself. Our recommended treatment strategy involves two major approaches:

- Promote autophagy to help rid the cells of the spike protein • Use interventions that limit the toxicity/pathogenicity of the spike protein

normalize the immune system rather than the use of immunosuppressant drugs, which may make the condition worse.

We recommend the use of immune-modulating agents and interventions to dampen and

Although we have listed suggested therapies below, we strongly suggest that, before initiating any of the below therapeutics, all patients and providers closely review the more detailed and comprehensive document — "An Approach to Managing Post-Vaccine Syndrome" — for information regarding dosing, cautions, contraindications, and other important details.

## First Line Therapies

(Not symptom specific; listed in order of importance)

- Intermittent daily fasting or periodic daily fasts Ivermectin
- Moderating physical activity
- Low-dose naltrexone
- Aspirin
- Nattokinase
- Melatonin
- Magnesium Methylene blue
- Sunlight and Photobiomodulation
- Resveratrol

# Probiotics/Prebiotics/Adjunctive/Second-Line Therapies

(Listed in order of importance)

- Vitamin D (with Vitamin K2)
- N-acetyl cysteine
- Cardio Miracle™ and L-arginine/L-citrulline supplements • Omega-3 fatty acids

• Intravenous Vitamin C

Non-invasive brain stimulation

- Sildenafil (with or without L-arginine- L-citrulline) • Nigella sativa
- Vitamin C • Spermidine
- Behavioral modification, relaxation therapy, mindfulness therapy, and psychological support

# Third Line Therapies

- Hyperbaric oxygen therapy
- Low Magnitude Mechanical Stimulation "Mitochondrial energy optimizer"

A note about anesthesia and surgery:

 Hydroxychloroquine Low-dose corticosteroid

Patients should notify their anesthesia team if using the following medications and/or nutraceuticals, as they can increase the risk of Serotonin syndrome (SS) with opioid administration: Methylene blue Curcumin Nigella Sativa Selective Serotonin Reuptake Inhibitors (SSRIs)

About Ivermectin Ivermectin is a well-known, FDA-approved drug that has been used successfully around the

world for more than four decades. One of the safest drugs known, it is on the WHO's list of essential medicines, has been given over 3.7 billion times, and won the Nobel Prize for its global and historic impacts in eradicating endemic parasitic infections in many parts of the world. Review the totality of supporting evidence for ivermectin in COVID-19.

It is likely that ivermectin and intermittent fasting act synergistically to rid the body of the spike protein.

It appears that vaccine-injured patients can be grouped into two categories: i) ivermectin

more difficult to treat and require more aggressive therapy. For ivermectin responders, prolonged and chronic daily treatment is often necessary to support their recovery. In many, if the daily ivermectin is discontinued worsening symptoms

responders and ii) ivermectin non-responders. This distinction is important, as the latter are

Ivermectin is best taken with or just following a meal, for greater absorption.

### This protocol is solely for educational purposes regarding potentially beneficial therapies for COVID-19. Never disregard professional medical advice because of something you have read

**DISCLAIMER** 

often recur within days.

on our website and releases. This protocol is not intended to be a substitute for professional medical advice, diagnosis, or treatment with regard to any patient. Treatment for an individual patient should rely on the judgement of a physician or other qualified health provider. Always seek their advice with any questions you may have regarding your health or medical condition. Please note our full disclaimer at: <a href="https://www.flccc.net/disclaimer">www.flccc.net/disclaimer</a> Download I-RECOVER Post-Vaccine Download I-RECOVER Post-Vaccine

Email Address\*

Last Name\*

Summary

Protocol

stay connected

SUPPORT US We are a 100% donor-supported 501(c)(3) non-profit organization. Our work would not be possible without you! Help

FLCCC®

ALLIANCE

**GIVE NOW** SHOP

us expand our reach and share life-saving research.

First Name\*

2001 L St. NW Suite 500 Washington, DC 20036

FLCCC Alliance

**CONTACT US** 

**MAILING ADDRESS** 

MEDIA press@flccc.net

> clinical@flccc.net **GENERAL**

**CLINICAL SUPPORT** 

support@flccc.net

#### > Protocols > Providers > Physicians

> About the FLCCC

**SHORTCUTS** 

SIGN ME UP

- Testimonials Latest Updates
- Newsletter Signup
- **FOLLOW US** f 💆 💿 🛠 🛊 🗮 🔘

**DISCLAIMER** 

©2020–2023 All Rights Reserved FLCCC Alliance. The information contained or presented on this website is for educational purposes only. Information on this site is NOT intended to serve as a substitute for diagnosis, treatment, or advice from a qualified, licensed medical professional. The facts presented are offered as information only in order to empower you – our protocol is not medical advice – and in no way should anyone infer that we, even though we are physicians, or anyone appearing in any content on this website are practicing medicine, it is for educational purposes only. Any treatment protocol you undertake should be discussed with your physician or other licensed medical professional. Seek the advice of a medical professional for proper application of ANY material on this site or our program to your specific situation. NEVER stop or change your medications without consulting your physician. If you are having an emergency contact your emergency services: in the USA that's 911. Please read our complete disclaimers.